

A Patient's Guide to **Osteochondritis Dissecans of the Knee**



©MMG 2003



Spine Patient Society™

Spine Patient Society
An educational and support community
for Spine Patients.
Philadelphia, PA 19147
Phone: ____-____-____ Fax: 251-352-6868

Spine Patient Society™ **Spine Patient Society**



The Spine Patient Society, a charitable, tax-exempt, 501(c)(3) organization, is dedicated to empowering patients with spine disorders through advocacy, education and global social support.

Our mission is to support all Spine Patients regardless of a chosen treatment modality, as we realize no two Spine Patients are alike. We believe that Spine Patients should not only support one another, but that they should also have access to professionally written, peer-reviewed educational materials. Our Society inspires Spine Patients to be proactive in their medical care and to become knowledgeable about their spinal disorders.



Spine Patient Society™

Spine Patient Society
An educational and support community
for Spine Patients.
Philadelphia, PA 19147
Phone: ____-____-____ Fax: 251-352-6868
justin@spinepatientsociety.org
www.SpinePatientSociety.org



Introduction


Osteochondritis dissecans (OCD) is a problem that affects the knee, mostly at the end of the big bone of the thigh (the *femur*). A joint surface damaged by OCD doesn't heal naturally. Even with surgery, OCD usually leads to future joint problems, including degenerative arthritis and osteoarthritis.

This guide will help you understand


- where in the knee the condition develops
- how doctors diagnose the problem
- what treatment options are available

Anatomy


What part of the knee is affected?



OCD mostly affects the *femoral condyles* of the knee. The femoral condyle is the rounded end of the lower thighbone, or *femur*. Each knee has two femoral condyles, referred to as the *medial femoral condyle* (on the inside of the knee) and the *lateral femoral condyle* (on the outside). Like most joint surfaces, the femoral condyles are covered in *articular cartilage*. Articular cartilage is a smooth, rubbery covering that allows the bones of a joint to slide smoothly against one another.



The problem occurs where the cartilage of the knee attaches to the bone underneath. The area of bone just under the cartilage surface is injured, leading to damage to the blood vessels of the bone. Without blood flow, the area of damaged bone actually dies. This area of dead bone can be seen on an X-ray and is sometimes referred to as the *osteochondritis lesion*.



The lesions usually occur in the part of the joint that holds most of the body's weight. This means that the problem area is under constant stress and doesn't get time to heal. It also means that the lesions cause pain and problems when walking and putting weight on the knee. It is more common for the lesions to occur on the medial femoral condyle, because the inside of the knee bears more weight.

Causes

How does the condition develop?

Juvenile Osteochondritis Dissecans

Children as young as nine or ten can develop this condition. But the disease behaves much differently in children and for this reason is given a separate name, *juvenile osteochondritis dissecans* (JOCD), meaning osteochondritis dissecans of children.

OCD and JOCD cause the same kind of damage to the knee, but they are separate diseases. In the child who is still growing, the problem is much more likely to heal itself. In the adult, the bones are not growing. For this reason, the treatment and prognosis of OCD and JOCD can be very different.

Many doctors think that JOCD is caused by repeated stress to the bone. Most young people with JOCD have been involved in competitive sports since they were very young. A heavy schedule of training and competing can stress the femur in a way that leads to JOCD. In some cases, other muscle or bone problems can cause extra stress and contribute to JOCD.

Osteochondritis Dissecans

Sometimes JOCD is not treated or does not heal completely. When this happens, JOCD develops into OCD. OCD can occur any time from early adulthood on, but most patients are adults under age 50. The cases of OCD that are first diagnosed in early adulthood probably began as JOCD. When a person gets OCD later in life, it is probably a brand new problem.

Doctors aren't sure what causes OCD. There is less of a link between strenuous, repetitive use and OCD. Many people who develop OCD don't have any particular risk factors.

Because OCD leads to damage to the surface of the joint, the condition can lead to problems with bone degeneration and osteoarthritis. The damage to the joint surface affects the way that the joint works. Like a machine that is out of balance, over time this imbalance can lead to abnormal wear and tear on the joint. This is one cause of degenerative arthritis and osteoarthritis.

Symptoms

What do OCD and JOCD feel like?


OCD and JOCD cause the same symptoms. The symptoms start out mild and grow worse with time. Both problems usually start with a mild aching pain. Moving the knee becomes painful, and it may be swollen and sore to the touch. Eventually, there is too much pain to put full weight on that knee. These symptoms are fairly common in athletes. They are similar to the symptoms of sprains, strains, and other knee problems.

As the condition becomes worse, the area of bone that is affected may collapse, causing a notch to form in the smooth joint surface. The cartilage over this dead section of bone (the lesion) may become damaged. This can cause a snapping or catching feeling as the knee joint moves across the notched area. In some cases the dead area of bone may actually become detached from the rest of the femur, forming what is called a *loose body*. This loose body may float around inside of the knee joint. The knee may catch or lock when it is moved if the loose body gets in the way.

Diagnosis

How do doctors identify this problem?

Your doctor will ask many questions about your medical history. You will be asked about



your current symptoms and about other knee or joint problems you have had in the past. Your doctor will then examine the painful knee by feeling it and moving it. You may be asked to walk, move, or stretch your knee. This may hurt, but it is important that your doctor knows exactly where and when your knee hurts.

Your doctor will probably order an X-ray of your knee. Most OCD lesions will show up on an X-ray of the knee. If not, your doctor may suggest a *bone scan*.

A bone scan involves injecting a special type of dye into the blood stream and then taking pictures of the bones with a special camera. This camera is similar to a Geiger counter and can pick up very small amounts of radiation. The dye that is injected is a very weak radioactive chemical. It attaches itself to areas of bone that are undergoing rapid changes, such as a healing fracture. A bone scan is the best way to see the lesions in the very early stages.

Your doctor may want to do other imaging tests, such as *magnetic resonance imaging* (MRI). The MRI machine uses magnetic waves rather than X-rays to show the soft tissues of the body. With this machine, doctors are able to create pictures that look like slices of the knee and see the anatomy, and any injuries, very clearly. These tests may help determine the extent of damage from OCD

and JOCD, and they also help rule out other problems.

Treatment

How do doctors treat the condition?

Many cases of JOCD can be completely healed with careful treatment. OCD will probably never completely heal, but it can be treated. There are two methods of treating JOCD: nonsurgical treatment to help the lesions heal, and surgery. Surgery is usually the only effective treatment for OCD.

Nonsurgical Treatment

Nonsurgical treatments help in about half the cases of JOCD. The goal is to help the lesions heal before growth stops in the thighbone. Even if imaging tests show that growth has already stopped, it is usually worth trying nonsurgical treatments. When these treatments work, the knee seems as good as new, and the JOCD doesn't seem to lead to arthritis.

Nonsurgical treatment of JOCD can take from 10 to 18 months. During that time, it is crucial to stop doing everything that causes pain to the knee. This means stopping exercise and sports. It may require using crutches or wearing a cast for a couple of months when symptoms are present. As knee symptoms ease, exercises can be started that don't involve placing weight through your foot. The exercises should be done carefully and should not cause any pain. Patients often work with physical therapists to develop an exercise program.

Your doctor may want to see what is happening in the knee and may suggest additional tests if your symptoms change. This may include new x-rays, MRI scan or a bone scan if your symptoms warrant additional testing. Even in JOCD, surgery may eventually be required. When the lesion has become so bad that it detaches totally or partially from the bone, nonsurgical treatment will not work. Even with the treatment, some patients

continue to have symptoms or their bone scans show signs that the damage is getting worse.

Some patients who are too near the end of bone growth may not benefit with nonsurgical treatment. When these problems develop, your surgeon may suggest surgery.

Surgery

If the lesion becomes totally or partially detached, surgery is needed to remove the loose body or to fix it in place. Your surgeon will need to gather lots of information about your knee and your problem before surgery.

This may require additional bone scans, X-rays, or MRIs. Your surgeon may also use an *arthroscope*, a tiny camera inserted into the knee to look at your knee before doing surgery to fix the problem. These tests are important because your surgeon needs to know the exact location and the size of the lesion to determine what kind of surgery will work best.

Arthroscopic Method

In some cases, your surgeon will be able to use the arthroscope to do the surgery. If the arthroscope can be used, the procedure requires smaller incisions than for an open surgery. This may reduce the time needed before the knee can be moved and exercised.


Open Method

Open surgery is needed when your surgeon can't get a picture of the entire lesion, when it is unclear how the fragment would best fit into the bone, or when it would be too difficult to replace the fragment using the arthroscope. Open surgery usually requires larger incisions than arthroscopic surgery to allow the surgeon to see into the knee and perform the operation.

Fragment Repair

If the loose bone fragment is in a weight-bearing area of your bone, your surgeon will try to reattach it if at all possible. Your surgeon may use tiny metal pins or screws to hold the

fragment in place. This sometimes proves difficult. The damaged fragment often doesn't fit perfectly into the bone anymore. And the bone around the fragment has often changed in ways that mean your surgeon will need to rebuild it.



Despite the difficulties, reattaching the fragment generally results in much better knee function than removing it. Your knee will not be as good as new, but a careful plan of exercise and follow-up care can help you use your knee again without pain.

Allograft Transplant

In rare cases, the lesion must be removed from a weight-bearing area. Your surgeon may try to fill in the hole using an *allograft*. An allograft is an actual transplant of bone and cartilage from a donor into your knee. The bone is usually obtained from a bone and tissue bank.

In this case, bone material is transplanted into the hole left in the bone. Allografts have risks, including graft rejection and infection. But they can be very successful in returning function to the knee.

Osteochondral Autograft

An *autograft* is a procedure for grafting tissue from the patient's own body. The place where the graft is taken is called the *donor site*. In

this case, surgeons graft a small amount of bone (*osteo*) and cartilage (*chondral*) from the donor site to put into the lesion. Usually, the donor site for this procedure is on the joint surface of the injured knee. Surgeons are careful to take the graft from a spot that won't cause a lot of problems, usually on the top and outside border of the knee cartilage. Even then, people sometimes end up with problems around the donor site. Surgeons have gotten good results with this surgery, but it is challenging to contour the graft to be just the same shape as the covering of the joint.

Autologous Chondrocyte Implantation

A new technology called *autologous chondrocyte implantation* is currently being developed. It involves using cartilage cells (*chondrocytes*) to help regenerate articular cartilage. This technology looks promising for treating JOCD and OCD but is still very much experimental.

Rehabilitation

What should I expect after treatment?

Nonsurgical Rehabilitation

The goal of nonsurgical rehabilitation is to help you learn ways to protect the injured area of cartilage while improving knee motion and strength. You may be advised to avoid heavy sport or work activities for up to eight weeks. Doing exercises in a pool can help you stay limber and fit while protecting the knee during this period.

Your doctor may have you work with a physical therapist for four to six weeks. Range-of-motion and stretching exercises are used to improve knee motion. Your therapist may issue shock-absorbing shoe insoles to reduce impact and protect your knee joint. You will also be shown strengthening exercises for the hip and knee to help steady the knee and give it additional protection from shock and stress.

After Surgery

If you have surgery, your surgeon may have you use a *continuous passive motion* (CPM) machine after surgery to help the knee begin to move and to alleviate joint stiffness.

With the exception of arthroscopic removal of a loose body, patients are instructed to avoid putting too much weight on their foot when standing or walking for up to six weeks. This gives the area time to heal. Weight bearing is usually restricted for up to four months after transplant procedures.

Patients are strongly advised to follow the recommendations about how much weight is safe. They may require a walker or pair of crutches for up to six weeks to avoid putting too much pressure on the joint when they are up and about.

Many surgeons will have their patients take part in formal physical therapy after knee surgery for osteochondritis lesions. The first few physical therapy treatments are designed to help control the pain and swelling from the surgery. Physical therapists will also work with patients to make sure they are only putting a safe amount of weight on the affected leg.

Exercises are chosen to help improve knee motion and to get the muscles toned and active again. At first, emphasis is placed on exercising the knee in positions and movements that don't strain the healing part of the cartilage. As the program evolves, more challenging exercises are chosen to safely advance the knee's strength and function.

Ideally, patients will be able to resume their previous lifestyle activities. Some patients may be encouraged to modify their activity choices, especially if an allograft was used.

The therapist's goal is to help you keep your pain under control, ensure safe weight bearing, and improve your strength and range of

motion. When you are well under way, regular visits to your therapist's office will end. The therapist will continue to be a resource, but you will be in charge of doing your exercises as part of an ongoing home program.

Notes