

A Patient's Guide to **Arthritis of the Thumb (CMC Joint)**



©MMG 2003

 **Northwestern**
Medicine

Orthopaedics

Northwestern Medicine Orthopaedics
27650 Ferry Road
Suite 100
Warrenville, IL 60555
Phone: 630.225.2663

Northwestern Medicine Orthopaedics


Northwestern Medicine Orthopaedics is pleased to provide this comprehensive library of orthopaedic and musculoskeletal patient information.

Simply click a topic of interest to access resourceful and helpful information.

These educational and easy-to-understand materials are always available for you on rmg.nm.org/orthopaedics.



Northwestern Medicine Orthopaedics
27650 Ferry Road
Suite 100
Warrenville, IL 60555
Phone: 630.225.2663
<http://www.oaortho.com>



Introduction


When you stop to think about how much you use your thumbs, it's easy to see why the joint where the thumb attaches to the hand can suffer from wear and tear. This joint is designed to give the thumb its rather large range of motion, but the tradeoff is that the joint suffers a lot of stress over the years. This can lead to painful osteoarthritis of this joint that may require surgical treatment as the arthritis progresses.

This guide will help you understand


- how arthritis of the thumb develops
- how it is diagnosed
- what can be done for the condition

Anatomy


Where is the CMC joint, and what does it do?



The CMC joint (an abbreviation for *carpo-metacarpal joint*) of the thumb is where the *metacarpal* bone of the thumb attaches to the *trapezium* bone of the wrist. This joint is sometimes referred to as the *basal joint* of the thumb. The CMC is the joint that allows you to move your thumb into your palm, a motion called *opposition*.



Several ligaments hold the CMC joint together. These ligaments can be injured, such as when you sprain your thumb. The joint surfaces are covered with a material called *articular cartilage*. This material is the slick, spongy covering that allows one side of a joint to slide against the other joint surface easily.




Causes


What causes arthritis of the thumb?

Arthritis is a condition in which a joint becomes *inflamed* (red, swollen, hot, and

painful). **Degenerative arthritis** is a condition in which a joint wears out, usually slowly over a period of many years. Doctors sometimes also describe this same condition as *degenerative arthrosis*. It is also called *osteoarthritis*.



Injury to a joint, such as a bad sprain or fracture, can cause damage to the articular cartilage. An injury to the CMC joint of the thumb, even if it does not injure the articular cartilage directly, can alter how the joint works. After a fracture of the thumb metacarpal, the bone fragments may heal in slightly different positions. The joints may then line up differently. This is also true when the ligaments around the CMC joint are damaged by a sprain. When an injury results in a change in the way the joint moves, the injury may increase the forces on the articular cartilage surfaces. This is similar to any mechanical device or machinery. If the mechanism is out of balance, it tends to wear out faster.




Over many years this imbalance in the joint mechanics can lead to damage on the articular surface. Since articular cartilage cannot heal itself very well, the damage adds up. Eventually, the joint is no longer able to compensate for the increasing damage, and it begins to hurt. Damage has occurred well before the pain begins.

Symptoms

What does arthritis of the thumb feel like?


Pain is the main problem with degenerative arthritis of any joint. This pain occurs at first only related to activity. Usually, once the activity gets underway there is not much pain, but after resting for several minutes the pain and stiffness increase. Later, when the condition worsens, pain may be present even at rest. The most noticeable problem with CMC joint arthritis is that it becomes difficult to grip anything. It causes a sharp pain at the **base of the thumb** in the thick part of the heel of the hand.



When the articular cartilage starts to wear off the joint surface, the joint may make a squeaking sound when moved. Doctors refer to this sound as *crepitus*. The joint often becomes stiff and begins to lose motion. Moving the thumb away from the palm may become difficult. This is referred to as a *contracture*.

Osteoarthritis may cause the CMC joint of the thumb to loosen and to bend back too far (*hyperextension*). If the middle thumb joint (MCP joint) becomes flexed and the furthest thumb joint also becomes hyperextended, the deformity is named a **thumb swan neck deformity**.

mity. A similar finger deformity sometimes occurs in people with finger arthritis.



Diagnosis

How do doctors identify this problem?


The diagnosis of CMC joint arthritis of the thumb begins with your doctor taking a detailed history of the problem. Specifics about any injuries that may have occurred to the hand are important because they may suggest other reasons why the condition exists.

Following the history, the doctor will examine your hand and possibly other joints in your body. The doctor will need to see how the motion of the CMC joint has been affected.

X-rays will be taken to see how much the joint is damaged. This test usually determines



how bad the degenerative arthritis has become. How much articular cartilage remains in the joint can be estimated with the X-rays.



Treatment

What can be done for CMC joint arthritis?

The treatment of degenerative arthritis of the CMC joint of the thumb can be divided into the nonsurgical means to control the symptoms and the surgical procedures that are available to treat the condition. Surgery is usually not considered until the symptoms have become impossible to control without it.

Nonsurgical Treatment

Treatment usually begins when the CMC joint first becomes painful. This may only occur with heavy use and may simply require mild anti-inflammatory medications, such as aspirin or ibuprofen. Reducing the activity, or changing from occupations that require heavy repetitive gripping with the hand, may be necessary to help control the symptoms.

Rehabilitation services, such as physical and occupational therapy, have a critical role in the nonoperative treatment plan for CMC joint arthritis. A primary goal is to help you learn how to control symptoms and maximize the health of your thumb. You'll learn ways to calm pain and symptoms, which might include the use of rest, heat, or topical rubs.

A brace or splint may also be prescribed to support the thumb. These devices are designed

to help reduce pain, prevent deformity, or keep a thumb deformity from getting worse. A *thumb stabilizer* is a type of thumb splint that is often custom-made of heat-moldable plastic. It is designed to fit the forearm, wrist, and thumb. Patients with CMC joint arthritis usually only wear the splint at night and when the joint is flared up. It should also be worn to protect the thumb during heavy or repeated hand and thumb activities.

Range-of-motion and stretching exercises are prescribed to improve your thumb motion. Strengthening exercises for the arm and hand help steady the hand and protect the thumb joint from shock and stress. Your therapist will go over tips on how you can get your tasks done with less strain on the joint.

An **injection of cortisone** into the joint can give temporary relief. Cortisone is a very powerful anti-inflammatory medication. When injected into the joint itself, it can help relieve the pain. Pain relief is temporary and usually only lasts several weeks to months. There is a small risk of infection with any injection into the joint, and cortisone injections are no exception.




Surgery

The surgical treatment for arthritis of the CMC joint includes several options. At one time, joint replacement with an artificial joint made with silicon was very popular. Problems with silicon implants in other parts of the body have led many surgeons to return to more traditional operations such as *fusion* and *excision arthroplasty* instead. Newer artificial joints are being developed, and in the future we may see more surgeons using them.

CMC Joint Fusion

A fusion, or *arthrodesis*, of any joint is designed to eliminate pain by allowing the bones that make up the joint to grow together, or fuse, into one solid bone. Fusions are used in many joints and were very common





before the invention of artificial joints for the replacement of arthritic joints. Even today, joint fusions are still commonly used in many different joints for treating the pain of arthritis.

A fusion of the CMC joint of the thumb is done quite often in younger people who need a strong grip or pinch more than they need the fine motion of the thumb. People who use their hands for heavy work will probably prefer a fusion over an arthroplasty (described below).


Artificial Joint Replacement (Arthroplasty)

Artificial joints are available for the CMC joint. These plastic or metal prostheses are used by some hand surgeons to replace the joint. The prosthesis acts as a spacer to fill the gap created when the arthritic surfaces of the two bones that make up the CMC joint are removed.



Excision Arthroplasty

The traditional operation for treating CMC joint arthritis is **excision arthroplasty**. This method has been used for many years and has withstood the test of time. The purpose of excision arthroplasty is to remove the arthritic joint surfaces of the CMC joint and replace them with a cushion of material that will



keep the bones separated. Most surgeons use a piece of tendon that has been rolled up and placed into the space created by removing the bone surfaces. During the healing phase after surgery, this tendon turns into tough scar tissue that forms a flexible connection between the bones, similar to a joint.

This operation is also combined with a reconstruction of the joint where tendons in the area are used to create a ligament sling between the metacarpal bone of the thumb and the carpal bone of the index finger. This helps hold the thumb in place and keeps the space between the bones from collapsing.

Rehabilitation

What should I expect following treatment?

Nonsurgical Rehabilitation

If you don't need surgery, range-of-motion exercises for the thumb should be started as pain eases. Your therapist will work with you to obtain or create a special thumb splint (mentioned earlier) when needed. The program advances to include strength exercises for the thumb and fingers. Dexterity and fine motor exercises are used to get your hand and thumb moving smoothly. You'll be given tips on keeping your symptoms controlled. You will probably progress to a home program within four to six weeks.

After Surgery

Your hand will be bandaged with a well-padded dressing and a thumb splint for support after surgery. Physical or occupational therapy sessions may be needed after surgery for up to eight weeks. The first few treatments are used to help control the pain and swelling after surgery. Some of the exercises you'll begin to do are to help strengthen and stabilize the muscles around the thumb joint. Other exercises are used to improve fine motor control and dexterity of your hand. You'll be given tips on ways to do your activities while avoiding extra strain on the thumb joint.

Notes