

A Patient's Guide to **Joint Injections for Arthritis**



©MMG 2005



Mackie Orthopaedics
Lower Level 2, Calvary Medical Centre
49 Augusta Road
Lenah Valley, Tas 7008
Phone: 61362281490 Fax: 61362281449



Mackie Orthopaedics



The information provided in these pages is licensed from an American surgical multimedia information provider. It represents general information on a variety of orthopaedic topics. The information may not always be entirely relevant to Australian assessments and management.


Dr Mackie does provide additional information to the majority of patients presenting for assessment and treatment of orthopaedic conditions. Questions arising from the content of these information pages may require a review consultation with Dr Mackie.

Dr Mackie has provided adult and paediatric orthopaedic care in Hobart since 2005. Services are provided to the Royal Hobart Hospital and all private hospitals. Most paediatric orthopaedic care is provided at the Royal Hobart Hospital due to the additional supports required.

Please contact Dr Mackie's office if you wish to arrange consultation or further information.



Mackie Orthopaedics
Lower Level 2, Calvary Medical Centre
49 Augusta Road
Lenah Valley, Tas 7008
Phone: 61362281490 Fax: 61362281449
clinic@mackie.net.au
<http://www.mackie.net.au>



Doctors recommend injections of *corticosteroids* (also commonly known as *cortisone*) for many arthritis patients. Cortisone is a powerful anti-inflammatory medication that can reduce joint inflammation. Because the medication is injected directly into the joint, the effects of the medication are concentrated on the painful joint. The injected cortisone can bring the inflammation in the joint under better control and decrease the swelling and pain.

These injections involve putting a needle directly into the joint. Through the needle, your doctor can remove excess *synovial fluid* (the lubricating fluid found in joints) and inject corticosteroid medication to help reduce the inflammation, pain, and swelling.

This process may sound risky. It is actually safe and fast. It involves little or no pain. And therapeutic injections have important benefits. They deliver the medicine to the exact spot that needs it. They also allow you to use lower and fewer doses of oral steroids, which are highly toxic.

Most doctors give only three to four injections per year in large, weight-bearing joints. This includes joints in your knee and hip. However,

patients with arthritis pain that cannot be controlled in other ways can get injections more often.

Complications

The most common side effect from injections is a temporary increase in pain and swelling. Rest, cold packs, and anti-inflammatory drugs help this pain go away within four to twenty-four hours. Studies have shown that about 6 percent of arthritis patients who receive injections in their joints experience this passing pain. It is probably caused by the body's reaction to the corticosteroid crystals in the medicine. If you have problems with pain and swelling after injections, your doctor may want to change the type of corticosteroid in your next injection.

Another fairly common complication is mild, temporary *flushing* (sudden redness of the skin) and agitation. Injections can also make diabetic symptoms worse.

There is a chance that the injection can introduce an infection into the joint. However, the odds of this are very slight. Studies show infections following injections happen from 1 in 1000 to 1 in 1600 times. Still, infections in the joint can be very serious. It is important to have an experienced professional perform the injection.

Some doctors and patients have wondered if the cartilage and other tissues of the joints are damaged by injections into the joint. Studies have not shown this to happen.

Notes